

On gradings of matrix algebras and descent theory

S. Caenepeel
Faculty of Applied Sciences
Free University of Brussels, VUB
B-1050 Brussels, Belgium

S. Dăscălescu and C. Năstăsescu*
Faculty of Mathematics
University of Bucharest
RO-70109 Bucharest 1, Romania

0 Introduction

The aim of this paper is to investigate group gradings of matrix algebras. Let k be a field and G be a group. A k -algebra A is called G -graded if $A = \bigoplus_{g \in G} A_g$, a direct sum of k -subspaces, such that $A_g A_h \subseteq A_{gh}$ for any $g, h \in G$. Several papers have been devoted to constructing and classifying group gradings on the matrix algebra $M_m(k)$. One can view the algebra $M_m(k)$ as a quotient of the path algebra of the quiver Γ , where Γ is the complete graph on m points. Certain gradings on $M_m(k)$ were constructed in [9], [10] from weight functions on Γ . These gradings were called *good gradings* in [6], where they were explained in an elementary way: they are the gradings for which all the matrix units e_{ij} are homogeneous elements. The name good gradings indicates that it is easy to understand and construct them. As we will see in this paper, good gradings play a central role in the classification of all gradings on the matrix algebra. There is one more way to present good gradings. If V is a G -graded vector space of dimension m , then there is a natural G -grading induced on $\text{End}(V)$ (denoted by $\text{END}(V)$), and this transfers to a grading of $M_m(k) \cong \text{End}(V)$. It turns out that this grading is a good grading, and moreover, any good grading can be obtained in this way. In Section 2 we prove that the gradings $\text{END}(V)$ and $\text{END}(W)$ are isomorphic if and only if the graded vector space W is a suspension of V . This result can be used to classify all good G -gradings on $M_m(k)$ by the orbits of the bi-action of the symmetric group S_m and G on the set G^m . We develop a combinatorial framework to use this for counting the good G -gradings in the case where $G = C_n$, the cyclic group of order n .

A fundamental result that we prove is that for an algebraically closed field k , any C_n -grading on $M_m(k)$ is isomorphic to a good grading. We give two proofs for this. The first one relies on a result of [6] concerned to graded Clifford theory, while the second one follows from an exact sequence of [1] concerned to graded Brauer groups. This fundamental result combined with the classification of the good C_n -gradings completes the classification of all C_n -gradings on $M_m(k)$ for an algebraically closed field k . Our approach to the case where k is arbitrary uses descent theory. Gradings of matrix algebras by the cyclic group C_n have been investigated in [3] with a different approach under the assumption that the field k contains a primitive n th root of unity. In this situation the group algebra kC_n is a selfdual Hopf algebra, and then a C_n -grading is the same to an action of C_n . The method of [3] consists of using the Skolem-Noether theorem and the Jordan form for presenting a C_n -grading. In this paper we do not need any assumption on the field. In Section 1 we present some preliminary results about descent theory. We use them to prove the main result of Section 3, stating that for a field k and a finite abelian group G , the forms of a certain (good) G -grading on $M_m(\bar{k})$ are in bijection to the isomorphism classes of Galois extensions of k with a certain Galois group. We are able to

*Research supported by the bilateral project "Hopf Algebras in Algebra, Topology, Geometry and Physics" of the Flemish and Romanian governments

give an explicit description of the grading corresponding to a Galois extension. We exemplify our theory by computing effectively the isomorphism types of C_2 -gradings on $M_2(k)$. This recovers and explains results of [3] in characteristic different from 2, and of [2] in characteristic 2. For proving the main result we need a graded version of the Rosenberg-Zelinsky exact sequence, and this has as a biproduct a graded version of the Skolem-Noether theorem.

1 Descent theory

In Section 3, we will use classical descent theory, and we briefly recall the basic results. Our main references are [12] and [14].

Let k be a commutative ring, and l a commutative faithfully flat k -algebra. Consider an “object” N (a module, an algebra, a graded algebra, ...) defined over l . Descent theory gives an answer to the following two problems:

- (i) *existence of forms*: does there exist an object M defined over k such that $M \otimes_k l \cong N$?
- (ii) *classification*: if such a k -form of N exists, classify all forms up to isomorphism.

The answer to the first question is basically the following (cf. [12, II.3.2]): M exists (as a k -module) if and only if there exists a *descent datum* for N , this is an $l \otimes_k l$ -isomorphism $u : l \otimes_k N \rightarrow N \otimes_k l$ such that $u_2 = u_3 \circ u_1$. Here $u_1 = I_l \otimes u : l \otimes_k l \otimes_k N \rightarrow l \otimes_k N \otimes_k l$, and u_2 and u_3 are defined in the same way.

If N is an l -algebra and u is an $l \otimes_k l$ -algebra map, then M is a k -algebra. Similar properties hold for other structures like F -graded algebras or F -module algebras (F being a group).

In principle M can be recovered from N , but this construction turns out to be too complicated in practical computations. However, if l is a Galois extension of k (in the sense of [7]), then we have a more explicit construction. Write $G = \text{Gal}(l/k)$, and take $\sigma \in G$. A map $f \in \text{End}_k(N)$ is called σ -semilinear if $f(an) = \sigma(a)f(n)$ for all $a \in l$ and $n \in N$. With these notations, we have the following result (cf. [12, III.5.1]):

Proposition 1.1 (Galois descent for modules) *Let l be a Galois extension of k with group G . $N \in l\text{-mod}$ descends to $M \in k\text{-mod}$ if and only if there exists a Galois descent datum for N , that is a group homomorphism*

$$\varphi : G \rightarrow \text{Aut}_k(N)$$

such that $\varphi(\sigma)$ is σ -semilinear for all $\sigma \in G$. The descended module M is the fixed module $M = N^G$, with the G -action on N induced by φ .

Again, similar results hold for other structures. For example, if $N = A$ is an l -algebra, then the descended module $M = B$ is a k -algebra if all the $\varphi(\sigma)$ are k -algebra maps.

Let us now look at the second problem. Assume that the l -module N descends to $M \in k\text{-mod}$. It is well-known (cf. [12, II.8]) that we have a bijection between the (pointed) set of isomorphism classes of k -modules M' that are isomorphic to $l \otimes_k M \cong N$ after base extension, and the Amitsur cohomology set $H^1(l/k, \text{Aut}_l(\bullet \otimes_k M))$:

$$\text{Mod}(l/k, M) \cong H^1(l/k, \text{Aut}_\bullet(\bullet \otimes_k M)) \quad (1)$$

We refer to [12] for the definition of Amitsur cohomology. We obtain similar statements about isomorphism classes of algebras, graded algebras, ... after we replace the automorphism group on the right hand side of (1) by the appropriate automorphism group, for example, for a k -algebra A , the set of algebra forms of $l \otimes_k A$ becomes isomorphic to

$$\text{Alg}(l/k, A) \cong H^1(l/k, \text{Aut}_\bullet^{\text{Alg}}(\bullet \otimes_k A)) \quad (2)$$

Taking inductive limits over all faithfully flat extensions l of k , we obtain a formula for all twisted forms of A as a k -algebra

$$\text{Alg}(k, A) \cong H^1(k_{\text{fl}}, \text{Aut}_{\bullet}^{\text{Alg}}(\bullet \otimes_k A)) \quad (3)$$

Here are some well-known explicit variations of (3), that will be used in the sequel. Taking $M = k^n$ in (1), we obtain the (generalized) Hilbert Theorem 90, after taking inductive limits:

$$\text{Mod}(l/k, k^n) \cong H^1(l/k, \text{Gl}_n) \quad (4)$$

If k is a field, then both sides of (4) are trivial, since there exists only one vector space of dimension n . For $n = 1$, we find

$$\text{Pic}(k) \cong H^1(l/k, \mathbb{G}_m) \quad (5)$$

where, as usual, $\mathbb{G}_m(k)$ denotes the group of invertible elements of k . Now let F be a finite group. The dual $Fk = \bigoplus_{\gamma \in F} kv_{\gamma}$ of the group algebra kF is an F -module algebra, the F -action is given by

$$\delta \cdot v_{\gamma} = v_{\gamma\delta^{-1}} \quad (6)$$

for all $\gamma, \delta \in F$. Fk is an F -Galois extension of k , and an F -module algebra form of Fk is again an F -Galois extension. Furthermore

$$\text{Aut}_F^{\text{alg}}(Fk) = F(k)$$

the group of all continuous maps from $\text{Spec}(k)$ to F , this is F to the power the number of connected components of k . We find the following formula for the set of commutative F -Galois extensions of k :

$$\text{Gal}(k, F) \cong H^1(k_{\text{fl}}, F) \quad (7)$$

The correspondence (1) is known in principle, but the construction is not very useful in practical situations. If l/k is Galois, then we have the following more explicit and practical description (see [5, Prop. 1.2] for the proof).

Proposition 1.2 *Let l be a Galois extension of the commutative ring k , with Galois group G , and consider a k -module M . The twisted form of $l \otimes_k M$ corresponding to a cocycle $\psi \in Z^1(G, \text{Aut}_l(l \otimes_k M))$ under the isomorphism*

$$\text{Mod}(l/k, M) \cong H^1(G, \text{Aut}_l(l \otimes_k M)) \quad (8)$$

is equal to $(l \otimes M)^G$, where the G -action on $l \otimes_k M$ is induced by the Galois descent datum $\varphi : G \rightarrow \text{End}_k(l \otimes_k M)$, with

$$\varphi(\sigma)(a \otimes m) = \psi(\sigma)(\sigma(a) \otimes m) = \sigma(a)(\psi(\sigma)(1 \otimes m)) \quad (9)$$

for all $a \in l$ and $m \in M$. Similar results hold for the variations of (8) (for algebras, coalgebras, Hopf algebras, ...).

2 Good gradings

Let k be a field, m a positive integer and G a group. We are interested to study G -gradings of the k -algebra of matrices $M_m(k)$. A special class of gradings consists of the good gradings. A grading is called good if all the matrix units e_{ij} are homogeneous elements. The class of good gradings is interesting since on one hand it is easy to construct them, and on the other hand any grading is isomorphic to a good one provided k , G and m satisfy certain conditions. This way to present good gradings from an interior point of view, by using certain elements of $M_m(k)$, has the advantage that one can easily generate the good gradings by assigning

arbitrary degrees to the elements $e_{12}, e_{23}, \dots, e_{m-1,m}$ (see [6, Proposition 2.1]). There is an alternative way to define good gradings, from an exterior point of view.

Let V be a G -graded k -vector space, i.e. $V = \bigoplus_{g \in G} V_g$, a direct sum of vector subspaces of V . For any $\sigma \in G$ we define the left σ -suspension $V(\sigma)$ of V as being the G -graded vector space which coincides with V as a vector space, and having the grading given by $V(\sigma)_g = V_{g\sigma}$ for any $g \in G$. Similarly, the right σ -suspension $(\sigma)V$ of V is just the space V with the grading $(\sigma)V_g = V_{\sigma g}$ for any $g \in G$. Obviously $((\sigma)V)(\tau) = (\sigma)(V(\tau))$ for any $\sigma, \tau \in G$, and we simply denote this by $(\sigma)V(\tau)$. We consider the endomorphism algebra $\text{End}(V)$ with map composition as multiplication. For any $\sigma \in G$ we denote by

$$\text{END}(V)_\sigma = \{f \in \text{End}(V) \mid f(V_g) \subseteq V_{\sigma g} \text{ for any } g \in G\}$$

Then the sum $\sum_{\sigma \in G} \text{END}(V)_\sigma$ is direct inside $\text{End}(V)$, and we denote by $\text{END}(V) = \bigoplus_{\sigma \in G} \text{END}(V)_\sigma$, which is a G -graded algebra (see [13]).

If V has dimension m , then $\text{END}(V) = \text{End}(V) \simeq M_m(k)$, and this provides a G -grading on $M_m(k)$ which turns out to be a good grading. In fact any good grading on $M_m(k)$ is isomorphic to $\text{END}(V)$ for some graded vector space V of dimension m (see [6, Proposition 1.2]), so from an exterior point of view we could define good gradings as gradings isomorphic to $\text{END}(V)$ for some V .

Our first aim is to classify the good G -gradings.

Theorem 2.1 *Let V and W be finite dimensional G -graded k -vector spaces. Then the graded algebras $\text{END}(V)$ and $\text{END}(W)$ are isomorphic if and only if there exists $\sigma \in G$ such that $W \simeq V(\sigma)$.*

Proof. Let $v_1, \dots, v_m$ be a basis of V consisting of homogeneous elements, say of degrees $g_1, \dots, g_m$. For any $1 \leq i, j \leq m$ define $E_{ij} \in \text{END}(V)$ by $E_{ij}(v_t) = \delta_{t,j} v_i$. Then E_{ij} is a homogeneous element of degree $g_i g_j^{-1}$ of $\text{END}(V)$, the set $(E_{ij})_{1 \leq i, j \leq m}$ is a basis of $\text{END}(V)$, and $E_{ij} E_{rs} = \delta_{j,r} E_{is}$ for any i, j, r, s . In particular $(E_{ii})_{1 \leq i \leq m}$ is a complete system of orthogonal idempotents of $\text{END}(V)$.

Let $u : \text{END}(V) \rightarrow \text{END}(W)$ be an isomorphism of G -graded algebras, and define $E'_{ij} = u(E_{ij})$ for any $1 \leq i, j \leq m$. Denote $Q_i = \text{Im}(E'_{ii})$, which is a graded vector subspace of W . Since $(E'_{ii})_{1 \leq i \leq m}$ is a complete system of orthogonal idempotents of $\text{END}(W)$, we have that $W = \bigoplus_{1 \leq i \leq m} Q_i$ and E'_{ii} acts as identity on Q_i for any i . Combined with the relation $E'_{ij} E'_{ji} = E'_{ii}$, this shows that E'_{ij} induces an isomorphism of degree $g_i g_j^{-1}$ from Q_i to Q_j . In particular $Q_j \simeq (g_1 g_j^{-1}) Q_1$ in $gr - k$ for any j . We obtain that $W \simeq \bigoplus_{1 \leq j \leq m} (g_1 g_j^{-1}) Q_1$, showing that Q_1 has dimension 1.

Repeating this argument for the identity isomorphism from $\text{END}(V)$ to $\text{END}(V)$, and denoting by $R_1 = \text{Im}(E_{11})$, we obtain that $V = \bigoplus_{1 \leq j \leq m} (g_1 g_j^{-1}) R_1$. Since Q_1 and R_1 are graded vector spaces of dimension 1, we have that $Q_1 \simeq R_1(\sigma)$ for some $\sigma \in G$. Hence $W \simeq \bigoplus_{1 \leq j \leq m} (g_1 g_j^{-1}) Q_1 \simeq \bigoplus_{1 \leq j \leq m} (g_1 g_j^{-1}) (R_1(\sigma)) \simeq V(\sigma)$. For the other way around, it is easy to see that $\text{END}(V) \simeq \text{END}(V(\sigma))$ as G -graded algebras. $\square$

The previous theorem can be used to classify all good G -gradings on the matrix algebra $M_m(k)$. Indeed, any such grading is of the form $\text{END}(V)$ for some G -graded vector space V of dimension m . To such a V we can associate an m -tuple $(g_1, \dots, g_m) \in G^m$ consisting of the degrees of the elements in a homogeneous basis of V . Conversely, to any such a m -tuple, one can associate a G -graded vector space of dimension m . Obviously, the G -graded vector space associated to a m -tuple coincides to the one associated to a permutation of the m -tuple. In fact Theorem 2.1 can be reformulated in terms of m -tuples as follows. If V and W are G -graded vector spaces of dimension m associated to the m -tuples $(g_1, \dots, g_m)$ and $(h_1, \dots, h_m)$, then $\text{END}(V) \simeq \text{END}(W)$ as G -graded algebras if and only if there exist $\sigma \in G$ and π a permutation of $\{1, \dots, m\}$ such that $h_i = g_{\pi(i)} \sigma$ for any $1 \leq i \leq m$. We have showed the following.

Corollary 2.2 *The good G -gradings of $M_m(k)$ are classified by the orbits of the biaction of the symmetric group S_m (from the left) and G (by translation from the right) on the set G^m .*

Classification of good gradings over a cyclic group

In the particular case where $G = C_n = \langle c \rangle$, a cyclic group with n elements, we are able to count the orbits of this biaction. Let $F_{n,m}$ be the set of all n -tuples $(k_0, k_1, \dots, k_{n-1})$ of non-negative integers with the property that $k_0 + k_1 + \dots + k_{n-1} = m$. It is a wellknown combinatorial fact that $F_{n,m}$ has $\binom{m+n-1}{n-1}$ elements. To any element $(g_1, \dots, g_m) \in G^m$ we can associate an element $(k_0, k_1, \dots, k_{n-1}) \in F_{n,m}$ such that k_i is the number of appearances of the element c^i in the m -tuple $(g_1, \dots, g_m)$ for any $1 \leq i \leq n$. By Corollary 2.2 we see that the number of orbits of the (S_m, G) -biaction on G^m is exactly the number of orbits of the left action by permutations of the subgroup $H = \langle \tau \rangle$ of S_n on the set $F_{n,m}$, where τ is the cyclic permutation $(1 \ 2 \ \dots \ n)$. This number of orbits is the one we will effectively compute.

If $\alpha = (k_0, k_1, \dots, k_{n-1}) \in F_{n,m}$ and d is a positive divisor of n , then $\tau^d \alpha = \alpha$ if and only if $k_{i+d} = k_i$ for any $0 \leq i \leq n-1-d$, i.e. the first d positions of α repeat $\frac{n}{d}$ times. In particular we must have that $\frac{n}{d}$ divides m . For any such d , let us denote by A_d the set of all $\alpha \in F_{n,m}$ stabilized by τ^d . Since for $\alpha \in A_d$ we have that $k_0 + k_1 + \dots + k_{d-1} = \frac{md}{n}$, we see that A_d has $\binom{\frac{md}{n} + d - 1}{d-1}$ elements.

Let $\mathcal{D}(n, m)$ be the set of all positive divisors d of n with the property that n/d divides m . If $d_1, d_2 \in \mathcal{D}(n, m)$, then $(d_1, d_2) \in \mathcal{D}(n, m)$. Indeed, since n divides $d_1 m$ and $d_2 m$, then n also divides $(d_1 m, d_2 m) = (d_1, d_2) m$, therefore $(d_1, d_2) \in \mathcal{D}(n, m)$. It follows that $\mathcal{D}(n, m)$ is a lattice with the order given by divisibility. For any $d \in \mathcal{D}(n, m)$ we denote by $\mathcal{D}(n, m, d)$ the set of all elements d' of $\mathcal{D}(n, m)$ which divide d , and by $\mathcal{D}_0(n, m, d)$ the set of all maximal elements of $\mathcal{D}(n, m, d)$. The following is immediate.

Lemma 2.3 *For any $d_1, d_2 \in \mathcal{D}(n, m)$ we have that $A_{d_1} \cap A_{d_2} = A_{(d_1, d_2)}$.*

The following describes the elements with the orbit of length d .

Lemma 2.4 *For any $d \in \mathcal{D}(n, m)$ denote by B_d the set of all the elements of $F_{n,m}$ having the orbit of length d . Then*

$$B_d = A_d - \bigcup_{d' \in \mathcal{D}(n, m, d)} A_{d'} = A_d - \bigcup_{d \in \mathcal{D}_0(n, m, d)} A_{d'}$$

Proof. The orbit of an element α has length d if and only if the stabilizer of α is a subgroup with $\frac{n}{d}$ elements of H , thus equal to $\langle \tau^d \rangle$. The result follows now from the definition of A_d . $\square$

Corollary 2.5 *Let $d \in \mathcal{D}(n, m)$ and $p_1, \dots, p_s$ all the distinct prime divisors of d such that $\frac{d}{p_1}, \dots, \frac{d}{p_s} \in \mathcal{D}(n, m)$. Then*

$$|B_d| = |A_d| + \sum_{1 \leq t \leq s} \sum_{1 \leq i_1 < \dots < i_t \leq s} (-1)^t |A_{\frac{d}{p_{i_1} \dots p_{i_t}}}|$$

and this is known taking into account the fact that for any d' we have that $|A_{d'}| = \binom{\frac{md'}{n} + d' - 1}{d' - 1}$.

Proof. We have that $\mathcal{D}_0(n, m, d) = \{\frac{d}{p_1}, \dots, \frac{d}{p_s}\}$. The result follows now from Lemma 2.4 and by applying the principle of inclusion and exclusion. $\square$

Theorem 2.6 *The number of isomorphism types of good C_n -gradings of the algebra $M_m(k)$ is*

$$\sum_{d \in \mathcal{D}(n, m)} \frac{1}{d} |B_d| = \sum_{d \in \mathcal{D}(n, m)} \frac{1}{d} (|A_d| - \bigcup_{d \in \mathcal{D}_0(n, m, d)} |A_{d'}|)$$

Proof. The number of isomorphism types of the good gradings is the number of orbits of the action of H on $F_{n,m}$. We have seen that if the orbit of an element $\alpha \in F_{n,m}$ has length d , then necessarily $d \in \mathcal{D}(n, m)$. The result follows since the number of orbits of length d is $\frac{1}{d}|B_d|$. $\square$

Example 2.7 Let $n = p^r$ with prime p and let m be a positive integer. We define q by $q = r$ in the case where n divides m , or q is the exponent of p in m in the case where n does not divide m . Then $\mathcal{D}(n, m) = \{p^i | r - q \leq i \leq r\}$ and the number of isomorphism types of good C_n -gradings on $M_m(k)$ is

$$\frac{1}{p^{r-q}} \binom{\frac{m}{p^q} + p^{r-q} - 1}{p^{r-q} - 1} + \sum_{r-q < i \leq r} \frac{1}{p^i} \left(\binom{\frac{m}{p^{r-i}} + p^i - 1}{p^i - 1} - \binom{\frac{m}{p^{r-i-1}} + p^{i-1} - 1}{p^{i-1} - 1} \right)$$

In particular, if $n = p$, then the number of isomorphism types of good C_p -gradings on $M_m(k)$ is $1 + \frac{1}{p} \left(\binom{m+p-1}{p-1} - 1 \right)$ if p divides m , and $\frac{1}{p} \binom{m+p-1}{p-1}$ if p does not divide m . This fact was proved in [3, Proposition 3.3] in the case where k contains a primitive p -th root of 1.

The algebraically closed field case

We show that in certain cases all gradings of the matrix algebras are isomorphic to good ones.

Theorem 2.8 *Let k be an algebraically closed field and m, n positive integers. Then any C_n -grading of the matrix algebra $M_m(k)$ is isomorphic to a good grading.*

Proof. Let $R = M_m(k)$ be the matrix algebra endowed with a certain C_n -grading, and pick $\Sigma \in R - \text{gr}$ a graded simple module. Then $\Delta = \text{End}_R(\Sigma) = \text{END}_R(\Sigma)$ is a C_n -graded algebra. Moreover, Δ is a crossed product when regarded as an algebra graded by the support of the C_n -grading of Δ . Thus $\Delta \simeq \Delta_1 \#_\sigma H$ for a cyclic group H and a cocycle σ . On the other hand $\Delta_1 = \text{End}_{R-\text{gr}}(\Sigma)$ is a finite field extension of k , so $\Delta_1 = k$. Hence Δ is a crossed product of Δ_1 , which is central in Δ , and the cyclic group H , so Δ is commutative. Since R is a semisimple algebra with precisely one isomorphism type of simple module, say S , we have that $\Sigma \simeq S^p$ as R -modules for some positive integer p . But then $\Delta \simeq \text{End}_R(S^p) \simeq M_p(k)$, and the commutativity of Δ shows that p must be 1. Then $\Sigma \simeq S$, so there exists a graded R -module which is simple as an R -module. By [6, Theorem 1.4], the grading is isomorphic to a good one. $\square$

3 Non-good gradings and descent theory

We have seen that all C_n -gradings on matrix algebras over an algebraically closed field are good. In this Section, we will classify C_n -gradings on matrix algebras over an arbitrary field, using descent. More precisely, we will discuss the following problem: given a finite abelian group G , and a G -graded k -vector space V , describe all gradings on $A = \text{End}_k(V)$ that become isomorphic to $\text{END}_{\bar{k}}(\bar{V})$ after a base extension, i.e. all gradings that are forms of the good grading $\text{END}_{\bar{k}}(\bar{V})$, where $\bar{k}$ stands for the algebraic closure of k and $\bar{V} = \bar{k} \otimes_k V$.

First, we introduce the following notation for a G -graded vector space V :

$$I(V) = \{\sigma \in G \mid V(\sigma) \cong V \text{ as graded vector spaces}\} \quad (10)$$

It is obvious that $I(V)$ is a subgroup of G . Also remark that $\sigma \in I(V)$ if and only if there exists an automorphism f of V that is homogenous of degree σ .

Lemma 3.1 *Let k be a field, G a finite abelian group, and V a G -graded vector space with $I(V) = H$. Then there exists a G -graded vector space W such that*

$$V \cong kH \otimes W \text{ and } I(W) = \{e\}$$

Proof. We choose coset representatives $e = \sigma_1, \sigma_2, \dots, \sigma_k$ of H in G , and let

$$W = \bigoplus_{i=1}^k V_{\sigma_i}$$

For all $\sigma \in H$, we can find an automorphism f_σ of V of degree σ , and this implies that V_{σ_i} and $V_{\sigma\sigma_i}$ are isomorphic, and have the same dimension. This can also be stated in the following way:

$$\bigoplus_{\sigma \in H\sigma_i} V_\sigma \cong kH \otimes V_{\sigma_i}$$

and it follows easily that $V \cong kH \otimes W$.

Now take $\sigma \in I(W)$. There exists an isomorphism $f : W \rightarrow W$ of degree σ , and $I_{kH} \otimes f : V \cong kH \otimes W \rightarrow V \cong kH \otimes W$ is also an isomorphism of degree σ , and this implies that $\sigma \in H$. If $\sigma \neq e$, then f sends V_{σ_i} to some V_{σ_j} , with $i \neq j$, implying that $\sigma_j = \sigma_i\sigma$, and contradicting $\sigma \neq e$. As a consequence, $\sigma = e$, and $I(W) = \{e\}$. $\square$

Now we fix a G -graded vector space V , and write $A = \text{END}_k(V)$. $\text{Aut}_{\text{gr}}(A)$ will denote the group of algebra automorphisms of degree e of A , and $\text{INN}(A)$ the subgroup consisting of inner algebra automorphisms induced by an isomorphism f of degree e of V . $\text{PC}(k, G)$ will be the group of grade preserving isomorphism classes of dimension one graded vector spaces. It is well-known and easy to show that $\text{PC}(k, G) \cong G$: the graded vector space $k(\sigma)$ represents the isomorphism class in $\text{PC}(k, G)$ corresponding to $\sigma \in G$.

The *Skolem-Noether Theorem* tells us that every automorphism of a matrix ring is inner. If we work over a commutative ring instead of a field, then this is no longer true, and the failure of the Skolem-Noether Theorem is measured by the *Picard group* of the ring, which is the set of isomorphism classes of rank one projective modules. In fact we have an exact sequence, known as the *Rosenberg-Zelinsky exact sequence*

$$0 \longrightarrow \text{INN}(A) \longrightarrow \text{Aut}(A) \xrightarrow{\varphi} \text{Pic}(k) \quad (11)$$

with $\varphi(\gamma)$ represented by $I = \{x \in A \mid x\gamma(a) = ax \text{ for all } a \in A\}$ and $\text{Im}(\varphi) = \{[I] \in \text{Pic}(k) \mid I \otimes V \cong V\}$ (see [12, IV.1.3] for the proof). Although we are interested in algebras over a field, we mention the commutative ring case because the graded version of the Picard group is not trivial, even if we work over a field: it is exactly the $\text{PC}(k, G)$ introduced above. We will need the following graded version of the Rosenberg-Zelinsky exact sequence (cf. [4, 13.6.1]):

Proposition 3.2 *Let G be a finite abelian group, V a G -graded vector space over a field k , and $A = \text{END}_k(V)$. Then we have an exact sequence*

$$0 \longrightarrow \text{INN}(A) \longrightarrow \text{Aut}_{\text{gr}}(A) \xrightarrow{\varphi} I(V) \longrightarrow 0$$

Proof. A straightforward adaption of the proof exhibited in [12, IV.1.3] gives an exact sequence

$$0 \longrightarrow \text{INN}(A) \longrightarrow \text{Aut}_{\text{gr}}(A) \xrightarrow{\varphi} \text{PC}(k, G)$$

with

$$\text{Im}(\varphi) = \{[I] \in \text{PC}(k, G) \mid I \otimes V \cong V\}$$

Now we have seen that $\text{PC}(k, G) \cong G$, and it is clear that $k(\sigma) \otimes V \cong V$ if and only if $\sigma \in I(V)$, and it follows that $\text{Im}(\varphi) \cong I(V)$. $\square$

Remark 3.3 The map $\varphi : \text{Aut}_{\text{gr}}(A) \rightarrow \text{PC}(k, G)$ can be described as follows: $\varphi(\gamma) = [I]$, with

$$I = \{x \in A \mid x\gamma(a) = ax \text{ for all } a \in A\}$$

Identifying $\text{PC}(k, G)$ and G , this means that $\varphi(\gamma) = \sigma$ if and only if γ is induced by an isomorphism of degree σ . This leads to the graded version of the Skolem-Noether Theorem: every isomorphism of a good graded matrix ring is induced by a homogeneous matrix.

Our next step is to introduce the category $\mathcal{G}_k$ of G -gradings on matrix rings over a field k . The objects are pairs

$$(V, \{A_\sigma \mid \sigma \in G\})$$

where V is a finite dimensional k -vector space, and

$$A = \text{End}_k(V) = \bigoplus_{\sigma \in G} A_\sigma$$

is a G -graded k -algebra. An (iso)morphism

$$(V, \{A_\sigma \mid \sigma \in G\}) \rightarrow (V', \{A'_\sigma \mid \sigma \in G\})$$

consists of an isomorphism of vector spaces $f : V \rightarrow V'$ such that the induced map $\text{End}(f) : A \rightarrow A'$ is a k -algebra isomorphism of degree e . The good gradings form a full subcategory $\mathcal{G}\mathcal{G}_k$ of $\mathcal{G}_k$. In the situation where $G = C_n$ and k is algebraically closed, all gradings are good, and the two categories coincide. We now want to apply (3), but with the category of k -algebras replaced by the category $\mathcal{G}_k$. First we have to compute the automorphisms of a good grading in the category $\mathcal{G}_k$.

Let V be a graded vector space, and $f : V \rightarrow V$ an isomorphism of the good grading $(V, \text{END}_k(V)) \in \mathcal{G}\mathcal{G}_k$. In Remark 3.3, we have seen that f is homogeneous of degree σ , for some $\sigma \in I(V) \subset G$. More precisely, we have the following commutative diagram:

$$\begin{array}{ccccccc}
& & 0 & & & & \\
& & \downarrow & & & & \\
& & \mathbb{G}_m(k) & & & 0 & \\
& & \downarrow & & & \downarrow & \\
0 & \longrightarrow & \text{Aut}_{\text{gr}}(V) & \longrightarrow & \text{Aut}_{\mathcal{G}_k}(V, \text{END}_k(V)) & \longrightarrow & I(V) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & \text{INN}(\text{END}_k(V)) & \longrightarrow & \text{Aut}_{\text{gr}}(\text{END}_k(V)) & \xrightarrow{\varphi} & I(V) \longrightarrow 0 \\
& & \downarrow & & & & \\
& & 0 & & & &
\end{array} \tag{12}$$

It follows from Proposition 3.2 and some diagram chasing that the rows and columns in the diagram are exact. Now we know from (3) (applied to the category of gradings), that the gradings on $\text{End}_k(V)$ that become isomorphic to the good grading $\text{END}_k(V)$ after a base extension are described by the first cohomology group $H^1(\bar{k}/k, \text{Aut}_{\mathcal{G}_{\bar{k}}}(\bar{V}, \text{END}_{\bar{k}}(\bar{V})))$, with $\bar{V} = \bar{k} \otimes V$. From the exactness of the second row in (12), it follows that we have an exact sequence of cohomology

$$H^1(\bar{k}/k, \text{Aut}_{\text{gr}}(\bar{V})) \rightarrow H^1(\bar{k}/k, \text{Aut}_{\mathcal{G}_{\bar{k}}}(\bar{V}, \text{END}_{\bar{k}}(\bar{V}))) \xrightarrow{\alpha} H^1(\bar{k}/k, I(\bar{V}))$$

Now we can write

$$\bar{V} = \bigoplus_{\sigma \in G} (k^{n(\sigma)})(\sigma)$$

where $n(\sigma)$ is the dimension of the part of degree σ of V ; this implies that

$$\text{Aut}_{\text{gr}}(V) = \bigoplus_{\sigma \in G} \text{GL}_{n(\sigma)}(\bar{k})$$

and $H^1(\bar{k}/k, \text{Aut}_{\text{gr}}(\bar{V}))$ becomes trivial, by the generalized Hilbert 90, cf. (4). This implies that the map α is injective. In fact we have the following:

Theorem 3.4 *Let k be a field, G a finite abelian group, and V a G -graded k -vector space. Then the map α described above is an isomorphism. Consequently, we have a bijective correspondence between*

- *isomorphism classes of gradings on $\text{End}_k(V)$ that become isomorphic to the grading $\text{END}_k(V)$ after base extension;*
- *isomorphism classes of Galois extensions l of k with Galois group $H = I(V)$.*

Proof. We have already seen that α is injective, so it suffices to show that α is surjective. The second statement then follows from the fact that the isomorphism classes of gradings are in bijective correspondence with $H^1(\bar{k}/k, \text{Aut}_{\mathcal{G}_{\bar{k}}}(\bar{V}, \text{END}_{\bar{k}}(\bar{V})))$, and isomorphism classes of H -Galois extensions are in bijective correspondence with $H^1(\bar{k}/k, H)$, cf. (7).

Consider a cocycle in $Z^1(\bar{k}/k, H)$. This cocycle corresponds to an H -Galois extension l of k , and l splits this cocycle, i.e. it is represented by some cocycle in

$$Z^1(H = \text{Gal}(l/k), H) = \text{Hom}(H, H)$$

and this cocycle is exactly the identity map $I_H : H \rightarrow H$. To prove that α is surjective, it suffices to find an inverse image in $H^1(H, \text{Aut}_{\mathcal{G}_l}(l \otimes V, \text{END}_l(l \otimes V)))$. Equivalently, we may look for the corresponding descent datum (see Proposition 1.2)

$$\varphi : H \rightarrow \text{Aut}_{\mathcal{G}_l}(l \otimes V, \text{END}_l(l \otimes V))$$

H is a finite abelian group, so we can write

$$H = \langle \sigma_1 \rangle \times \cdots \times \langle \sigma_n \rangle$$

with $\sigma_i^{m_i} = e$. We know φ once we know all the $\varphi(\sigma_i)$. Recall from Lemma 3.1 that

$$V \cong kH \otimes W$$

so

$$l \otimes V \cong l \otimes k \langle \sigma_1 \rangle \otimes \cdots \otimes k \langle \sigma_n \rangle \otimes W$$

We define $\varphi(\sigma_i) : l \otimes V \rightarrow l \otimes V$ as follows:

$$\varphi(\sigma_i) = \sigma_i \otimes I_{k\langle\sigma_1\rangle} \otimes \cdots \otimes \varphi_i \otimes \cdots \otimes I_{k\langle\sigma_n\rangle} \otimes I_W$$

with $\varphi_i : k\langle\sigma_i\rangle \rightarrow k\langle\sigma_i\rangle$ given by

$$\varphi_i(\sigma_i^l) = \sigma_i^{l+1}$$

i.e. the matrix of φ_i with respect to the basis $\{e, \sigma_i, \sigma_i^2, \dots, \sigma_i^{m_i-1}\}$ is

$$\begin{pmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \end{pmatrix}$$

Clearly $\varphi(\sigma_i)$ is σ_i -semilinear, and $\varphi(\sigma_i)^{m_i} = 1$, which means that φ is a descent datum. For all $\sigma \in H$, $\varphi(\sigma)$ is homogeneous of degree σ , and the same thing holds for the corresponding cocycle ψ . This means exactly that α maps ψ to the identity map I_H , proving that α is surjective. $\square$

Remark 3.5 Let l be an H -Galois extension of k . Let us give an explicit description of the grading corresponding to l on $\text{End}_k(kH)$. To this end, we first recall the definition of a Galois extension of a commutative ring (see [7, III.1.2]). Let l/k be an extension of commutative rings, and assume that H acts on l as a group of k -automorphisms. Now we consider the graded k -algebra

$$\Delta(l, H) = \bigoplus_{\sigma \in H} l u_\sigma$$

with multiplication

$$(a u_\sigma)(b u_\tau) = a \sigma(b) u_{\sigma\tau}$$

and H -grading

$$\deg(u_\sigma) = \sigma$$

We have a k -algebra map

$$j : \Delta(l, H) \rightarrow \text{End}_k(l) ; j(a u_\sigma)(b) = a \sigma(b)$$

and l is a Galois extension of k with group H if and only if $l^H = k$ and j is an isomorphism. If j is an isomorphism, then the grading on H induces a grading on $\text{End}_k(l)$, and this is the grading we are looking for.

Let us show that this grading is a form of the good grading on $\text{End}_k(kH)$. Since l is a Galois extension of k with group H , we have an isomorphism

$$m : l \otimes l \rightarrow Hl ; m(a \otimes b) = \sum_{\sigma \in H} a \sigma(b) v_\sigma$$

(see [7, III.1.2]). This means in fact that l is a form of the dual of the groupalgebra Hk , and it suffices to show that we have a graded isomorphism

$$f : \Delta(Hk, H) \rightarrow \text{End}_k(kH)$$

Let $\{e_{\sigma, \tau} \mid \sigma, \tau \in H\}$ be the canonical basis of $\text{End}_k(kH)$ consisting of elementary matrices with respect to the basis H of kH , i.e.

$$e_{\sigma, \tau} e_{\rho, \nu} = \delta_{\tau, \rho} e_{\sigma, \nu} \text{ and } \deg(e_{\sigma, \tau}) = \tau \sigma^{-1}$$

$\Delta(Hk, H)$ has k -basis $\{v_\sigma u_\tau \mid \sigma, \tau \in H\}$, with

$$(v_\sigma u_\tau)(v_\rho u_\nu) = \delta_{\sigma, \rho\tau^{-1}} v_\sigma u_{\tau\nu} \text{ and } \deg(v_\sigma u_\tau) = \tau$$

Now define

$$f(e_{\sigma, \tau}) = v_\sigma u_{\tau\sigma^{-1}}$$

It is straightforward to show that f is a grade preserving isomorphism of k -algebras.

Remark 3.6 Our results are closely related to an exact sequence of Beattie [1], which we will recall here in the particular situation that is of interest to us. Let k be a field, G a finite abelian group, and look at G -graded central simple algebras (in particular graded matrix algebras). Two such graded algebras A and B are called graded equivalent if there exist matrix rings $M_n(k)$ and $M_m(k)$ equipped with a good grading such that $A \otimes M_n(k)$ and $B \otimes M_m(k)$ are isomorphic as graded algebras. Equivalence classes of graded central simple algebras form a group under the operation induced by the tensor product, which we call the G -graded Brauer group of k , and we denote it by $BC(k, G)$. Beattie [1] proved that we have a split exact sequence

$$1 \rightarrow \text{Br}(k) \rightarrow BC(k, G) \rightarrow \text{Gal}'(k, G) \rightarrow 1 \quad (13)$$

Here $\text{Gal}'(k, G)$ consists of all isomorphism of (possibly noncommutative) Galois extensions of k with group G . The previously introduced $\text{Gal}(k, G)$ contains only commutative Galois extensions. If G is cyclic, then all Galois extensions are commutative. If $k = \bar{k}$ is algebraically closed, then $\text{Gal}(k, G)$ is trivial. In particular, it follows that $\text{Br}(\bar{k}) = BC(\bar{k}, C_n)$, implying that any C_n -grading on a matrix algebra over an algebraically closed field is good, providing an alternative proof of Theorem 2.8. The inverse image of a (commutative) Galois extension l of k in (13) is represented by the graded algebra constructed in Remark 3.5.

Example 3.7 In some cases, we can do the descent of the grading explicitly. Let k be a field of characteristic different from 2, and take $G = C_2 = \{e, \sigma\}$. We will look at forms of the nontrivial good grading on $M_2(k)$, given by

$$\deg(e_{11}) = \deg(e_{22}) = e ; \deg(e_{21}) = \deg(e_{12}) = \sigma$$

which is also the good grading $\text{END}_k(V)$, with $V = ke \oplus k\sigma$ with the obvious grading. From Theorem 3.4, we know that isomorphism classes of these gradings are given by the quadratic extensions $\text{Gal}(k, C_2) \cong k^*/(k^*)^2$. Recall that the quadratic extension corresponding to $\alpha \in k^*$ is

$$l = k[X]/(X^2 - \alpha)$$

Let $\sqrt{\alpha}$ be the element in l represented by X . The descent datum corresponding to l is

$$\sigma \otimes \varphi : l \otimes V \rightarrow l \otimes V$$

where the matrix of φ is

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

The induced map $\varphi = \text{End}(\varphi) : M_2(k) \rightarrow M_2(k)$ is given by

$$\varphi \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} d & c \\ b & a \end{pmatrix}$$

The descended graded matrix ring consists of the matrices in $M_2(l)$ that are fixed by $\sigma \otimes \varphi$, it is

$$A = \left\{ \begin{pmatrix} a & b \\ \sigma(a) & \sigma(b) \end{pmatrix} \mid a, b \in l \right\}$$

or

$$A = \left\{ \begin{pmatrix} a + b\sqrt{\alpha} & c + d\sqrt{\alpha} \\ c - d\sqrt{\alpha} & a - b\sqrt{\alpha} \end{pmatrix} \mid a, b, c, d \in l \right\}$$

A is the k -endomorphism ring of the descended module $\tilde{V}$, which consists of the elements in $l \otimes V$ fixed by $\sigma \otimes \varphi$:

$$\tilde{V} = \{(a + b\sqrt{\alpha})e + (a - b\sqrt{\alpha})\sigma \mid a, b \in k\}$$

A k -basis for $\tilde{V}$ is $\{u = e + \sigma, v = e - \sigma\}$.

Now we observe that

$$A_e = k \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \oplus k\sqrt{\alpha} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} ; A_\sigma = k \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus k\sqrt{\alpha} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Switching to the new basis $\{u, v\}$, we find the following grading on $B = M_2(k)$:

$$B_e = k \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \oplus k \begin{pmatrix} 0 & 1 \\ \alpha & 0 \end{pmatrix} ; B_\sigma = k \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus k \begin{pmatrix} 0 & 1 \\ -\alpha & 0 \end{pmatrix}$$

and we have recovered [3, Corollary 3.6]

Example 3.8 Now let k be a field of characteristic 2. Again we describe the grading forms on $M_2(k)$ with the same good grading as in Example 3.7. The difference with Example 3.7 is that the description of the isomorphism classes of quadratic extension is different. We now have

$$\text{Gal}(k, C_2) \cong k/\{x^2 - x \mid x \in k\}$$

The quadratic extension corresponding to $\alpha \in k$ is now

$$l = k[X]/(X^2 - X - \alpha)$$

We write x for the element in l represented by X . The action of C_2 on l is given

$$\sigma(x) = x + 1$$

Proceeding as in Example 3.7, we now find for the descended graded matrix ring

$$A = \left\{ \begin{pmatrix} a + bx & c + dx \\ c + d(x+1) & a + b(x+1) \end{pmatrix} \mid a, b, c, d \in k \right\}$$

The grading is now described by the formulas

$$A_e = k \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \oplus k \begin{pmatrix} x & 0 \\ 0 & x+1 \end{pmatrix} ; A_\sigma = k \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \oplus k \begin{pmatrix} 0 & x \\ x+1 & 0 \end{pmatrix}$$

and A is the k -endomorphism ring of the k -module

$$\tilde{V} = \{(a + bx)e + (a + b(x+1))\sigma \mid a, b \in k\}$$

Now we take the following k -basis of $\tilde{V}$:

$$\{u = e + \sigma, v = xe + (x+1)\sigma\}$$

If we write the k -endomorphisms of $\tilde{V}$ as matrices with respect to the basis $\{u, v\}$, then we find the following grading on $B = M_2(k)$:

$$B_e = k \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \oplus k \begin{pmatrix} 0 & \alpha \\ 1 & 1 \end{pmatrix} ; B_\sigma = k \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \oplus k \begin{pmatrix} 0 & \alpha \\ 1 & 0 \end{pmatrix} \quad (14)$$

As a consequence, we have now the list of all (different) isomorphism types of C_2 -graded algebra structures on $M_2(k)$, with k a field of characteristic 2, namely

- the trivial grading: all matrices have degree e ;
- the good grading $\deg(e_{11}) = \deg(e_{22}) = e$, $\deg(e_{12}) = \deg(e_{21}) = \sigma$;
- the bad gradings from (14), where α ranges over a system of representatives of the $\{x^2 - x \mid x \in k\}$ cosets of k different from $\{x^2 - x \mid x \in k\}$.

Using direct computations, these gradings have been described before in [6], and classified up to isomorphism in [2].

References

- [1] M. Beattie, A direct sum decomposition for the Brauer group of H -module algebras, *J. Algebra* **43** (1976), 686-693.
- [2] C. Boboc, Superalgebra structures on $M_2(k)$ in characteristic 2, *Comm. Algebra*, to appear.
- [3] C. Boboc and S. Dăscălescu, Gradings of matrix algebras by cyclic groups, *Comm. Algebra*, to appear.
- [4] S. Caenepeel, Brauer groups, Hopf algebras and Galois theory, Kluwer Academic Publ., Dordrecht, 1998.
- [5] S. Caenepeel, S. Dăscălescu and L. Le Bruyn, Forms of pointed Hopf algebras, *Manuscripta Math.* **100** (1999), 35-53.
- [6] S. Dăscălescu, B. Ion, C. Năstăsescu and J. Rios Montes, Group Gradings on Full Matrix Rings, *J. Algebra* **220** (1999), 709-728.
- [7] F. DeMeyer, E. Ingraham, Separable algebras over commutative rings, *Lecture Notes in Math.* **181**, Springer Verlag, Berlin, 1971.
- [8] J. L. Gomez Pardo and C. Năstăsescu, Graded Clifford theory and duality, *J. Algebra* **162** (1993), 28-45.
- [9] E. L. Green, Graphs with relations, covering and group-graded algebras, *Trans. Amer. Math. Soc.* **279** (1983), 297-310.
- [10] E. L. Green and E. N. Marcos, Graded quotients of path algebras: A local theory, *J. Pure Appl. Algebra* **93** (1994), 195-226.

- [11] M.A. Knus, Algebras graded by a group, in “Category theory, homology theory and their applications II”, P. Hilton (Ed.), *Lecture Notes in Math.* **92**, Springer Verlag, Berlin 1969, 117-133.
- [12] M. Knus and M. Ojanguren, “Théorie de la Descente et Algèbres d’Azumaya”, *Lect. Notes in Math.* **389**, Springer Verlag, Berlin, 1974.
- [13] C. Năstăsescu and F. Van Oystaeyen, Graded Ring Theory, North Holland, 1982.
- [14] J.P. Serre, “Cohomologie Galoisienne”, *Lect. Notes in Math.* **5**, Springer Verlag, Berlin, 1965.